

Dimension of Spaces of Automorphic Forms*

by

R.P. Langlands

I will first formulate a problem in the theory of group representations and show how to solve it; then I will discuss the relation of this problem to the theory of automorphic forms. Since there is no point in striving for maximum generality, I start with a connected semisimple group G with finite center. An irreducible unitary representation π of G on the Hilbert space H is said to be square-integrable if for one and hence, as one can show, every pair u and v of nonzero vectors in H the function $(\pi(g)u, v)$ is square-integrable on G . It is said to be integrable if for one such pair $(\pi(g)u, v)$ is integrable.

Suppose Γ is a discrete subgroup of G and $\Gamma \backslash G$ is compact. As was shown by Godement in an earlier lecture the representation π of the previous paragraph occurs a finite number of times, say $N(\pi)$, in the regular representation on $L^2(\Gamma \backslash G)$. The problem is first to find a closed formula for $N(\pi)$. The method which I will now describe of obtaining such a formula is valid only when π is actually integrable.

Square integrable representations are similar in some respects to representations of compact groups; in particular they satisfy a form of the Schur orthogonality relations. There is a constant d_π called the formal degree of π such that if u', v', u , and v belong to H then

$$\int_G (\pi(g)u', v') \overline{(\pi(g)u, v)} dg = d_\pi^{-1}(u', u)(v, v').$$

If u and v are such that $(\pi(g)u, v)$ is integrable and π' is unitary representation of G on H' which does not contain π , then

$$\int_G (\pi'(g)u', v') \overline{(\pi(g)u, v)} dg = 0$$

for all u', v' in H' .

* Appeared first in Proc. of Symp. in Pure Mathematics, vol IX, Amer. Math. Soc. (1966)

Let $L_i, 1 \leq i \leq N(\pi)$, be a family of mutually orthogonal invariant subspaces of $L^2(\Gamma \backslash G)$ which are such that the action of G on each of them is equivalent to π . Suppose that π does not occur in the orthogonal complement of

$$\sum_{i=1}^{N(\pi)} \oplus L_i.$$

If π is integrable there is a unit vector v in H such that $(\pi(g)v, v)$ is integrable. Let v_i be a unit vector in L_i corresponding to v under some equivalence between H and L_i . The orthogonality relations imply that the operator $\Phi \rightarrow \Phi'$ with

$$\begin{aligned} \Phi'(g) &= d_\pi \int_G \Phi(gh) \overline{(\pi(g)v, v)} dh, \\ &= \int_{\Gamma \backslash G} \Phi(g) \left\{ \sum_{\Gamma} \xi(g^{-1}\gamma h) \right\} dh, \end{aligned}$$

if $\xi(g) = d_\pi \overline{(\pi(g)v, v)}$, is an orthogonal projection on the space spanned by $v_1, \dots, v_{N(\pi)}$. For our purposes it may be assumed that v transforms according to a finite-dimensional representation of some maximal compact subgroup of G . Then the argument used by Borel in a previous lecture shows that

$$\sum_{\Gamma} \xi(g^{-1}\gamma h)$$

converges absolutely uniformly on compact subsets of $G \times G$. Hence $v_1, \dots, v_{N(\pi)}$ may be supposed continuous. As a consequence

$$\sum_{i=1}^{N(\pi)} v_i(g) \bar{v}_i(g) = \sum_{\Gamma} \xi(g^{-1}\gamma h).$$

Set $h = g$ and integrate over $\Gamma \backslash G$ to obtain

$$N(\pi) = \int_{\Gamma \backslash G} \sum_{\Gamma} \xi(g^{-1}\gamma g) dg.$$

The sum in the integrand may be rearranged at will. If Σ is a set of representatives for the conjugacy classes in Γ the integral on the right equals

$$\begin{aligned} \int_{\Gamma \backslash G} \sum_{\gamma \in \Sigma} \sum_{\delta \in \Gamma_\gamma \backslash \Gamma} \xi(g^{-1}\delta^{-1}\gamma\delta g) dg &= \sum_{\gamma \in \Sigma} \int_{\Gamma_\gamma \backslash G} \xi(g^{-1}\gamma g) dg \\ &= \sum_{\gamma \in \Sigma} \mu(\Gamma_\gamma \backslash G_\gamma) \int_{G_\gamma \backslash G} \xi(g^{-1}\gamma g) dg, \end{aligned}$$

if Γ_γ and G_γ are the centralizers of γ in Γ and G respectively. The equality of $N(\pi)$ and the final expression is of course a special case of a formula of Selberg and has been known for some time.

The problem of evaluating $\mu(\Gamma_\gamma \backslash G_\gamma)$, the volume of $\Gamma_\gamma \backslash G_\gamma$, has been discussed in the lectures on Tamagawa numbers. So we shall not worry about it now. Since $\Gamma \backslash G$ is compact every element of Γ is semisimple; thus our problem is to express the integral

$$\int_{G_\gamma \backslash G} \xi(g^{-1}\gamma g) dg$$

in elementary terms when γ is a semisimple element of G .

If π is a square-integrable representation of G on H , v is a vector in H which transforms according to a finite-dimensional representation of some maximal compact subgroup of G , and

$$\xi(g) = d_\pi(\overline{\pi(g)v}, v),$$

then a recent theorem of Harish-Chandra states that

$$(a) \quad \int_{G_\gamma \backslash G} \xi(g^{-1}\gamma g) dg$$

exists for γ semisimple and vanishes unless γ is elliptic, that is, belongs to some compact subgroup of G . Since Σ contains only a finite number of elliptic elements the sum in the expression for $N(\pi)$ is finite. We still require a closed expression for the integrals appearing in it.

Let K be a maximal compact subgroup of G . Since G has a square-integrable representation there is a Cartan subgroup T of G contained in K . It is enough to compute the integral (a) for γ in T . There is a limit formula of Harish-Chandra which allows one to compute its value at the singular elements once its values at the regular elements are known. Thus we need only evaluate it when γ is regular. It should be remarked that in this limit formula there is a constant which depends on the choice of Haar measure on G_γ . The exact relation of this constant to the choice of Haar measure has never been determined; until it is, our problem cannot be regarded as completely solved.

If γ is regular and the measure on G_γ is so normalized that the volume of G_γ is one, then

$$\int_{G_\gamma \backslash G} \xi(g^{-1}\gamma g) dg = \chi_\pi(\gamma^{-1})$$

if χ_π is the character of π . An explicit expression for the right-hand side has recently been obtained.

Let \mathfrak{h} be the Lie algebra of T ; choose an order on the roots of \mathfrak{h}_c ; and let Λ be a linear function on \mathfrak{h}_c such that $\Lambda + \rho, \rho = \frac{1}{2} \sum_{\alpha > 0} \alpha$, extends to a character of T . To each such Λ there is associated a square-integrable representation π_Λ and if $H \in \mathfrak{a}$

$$\chi_{\pi_\Lambda}(\exp H) = (-1)^m \epsilon(\Lambda) \sum_{\sigma \in W} \frac{\text{sgn} \sigma \exp(\sigma(\Lambda + \rho))(H)}{(\exp(\alpha(G)/2) - \exp(-\alpha(G)/2))} .$$

Here $m = \frac{1}{2} \dim G/K$, $\epsilon(\Lambda) = \text{sgn}(\prod_{\alpha > 0} (\Lambda + \rho, \alpha))$, and W is the Weyl group of K . Every square-integrable representation is equivalent to π_Λ for some Λ . However the values of Λ for which π_Λ is integrable are not yet known. For some special cases see [1] and [2].

The geometrical meaning of the numbers $N(\pi_\Lambda)$ is not yet completely clear. I would like to close this lecture with some suggestions as to what it might be. Since the evidence at present is rather meagre, they are only tentative. If $\mathfrak{g}_\mathbb{C}$ is the complexification of the Lie algebra of \mathfrak{g} , the elements of $\mathfrak{g}_\mathbb{C}$ may be regarded as left-invariant complex vector fields on G and G/T may be turned into a complex manifold in such a way that the space of antiholomorphic tangent vectors at $\bar{g} = gT$ is the image of $\mathfrak{n}_\mathbb{C}^-$ if $\mathfrak{n}_\mathbb{C}^-$ is the subalgebra of $\mathfrak{g}_\mathbb{C}$ generated by root vectors belonging to negative roots. Let V^* be the bundle of antihomomorphic cotangent vectors and introduce a G -invariant metric in V^* and hence in $\Lambda^q V^*$. Let B be the line bundle over G/T associated to the character $\xi(\exp H) = \exp(\Lambda(H))$ of T . If Γ is a discrete subgroup of G let $C^q(\Lambda, \Gamma)$ be the space of Γ -invariant cross-sections of $B \otimes \Lambda^q V^*$ which are square integrable over $\Gamma \backslash G/T$. There is a unique closed operator $\bar{\partial}$ from $C^q(\Lambda, \Gamma)$ to $C^{q+1}(\Lambda, \Gamma)$ whose domain contains the infinitely differentiable cross-sections of compact support on which $\bar{\partial}$ is to have its usual meaning and whose adjoint is defined on the infinitely differentiable cross-sections of $C^{q+1}(\Lambda, \Gamma)$ with compact support.

Set $C^q(\Lambda, \{1\}) = C^q(\Lambda)$. I expect, although I do not know how to prove it, that when $\Lambda + \rho$ is nonsingular the range of $\bar{\partial}$ is closed for every q . If this is so then the cohomology groups $H^1(\Lambda)$ will be Hilbert spaces on which G acts. Is it true that they vanish for all but one value of q , say $q = q_\Lambda$, and that the representation π'_Λ of G on $H^{q_\Lambda}(\Lambda)$ is equivalent to π_Λ ? The following theorem is a clue to the value of q_Λ .

Theorem (P. Griffiths) . Let a_1 be the number of noncompact positive roots for which $(\Lambda + \rho, \alpha) > 0$ and let a_2 be the number of compact positive roots for which $(\Lambda + \rho, \alpha) < 0$. There is a constant c such that if $|(\Lambda + \rho, \alpha)| > c$ for every simple root, $\Lambda \backslash G$ is compact, and Γ acts freely on G/T , then $H^q(\Lambda, \Gamma) = 0$ unless $q = a_1 + a_2$.

It is, I think, worthy of remark that if one assumes that $H^q(\Lambda) = \{0\}$ for $q \neq q_\Lambda = a_1 + a_2$, then a formal application of the Woods Hole fixed point formula shows that if γ is a regular element of T , then the value at γ of the character of π'_Λ is $\chi_{\pi_\Lambda}(\gamma)$. By the way, it is known that $H^0(\Lambda) = 0$ unless $q_\Lambda = 0$ and that if $q_\Lambda = 0$ the representation of G on $H^0(\Lambda)$ is in fact π_Λ .

Finally one will want to show that when π_Λ is integrable and $\Gamma \backslash G$ is compact the number $N(\pi_\Lambda)$ is equal to the dimension of $H^{q_\Lambda}(\Lambda, \Gamma)$. This can be done when $q_\Lambda = 0$; in this case $H^0(\Gamma, \Lambda)$ is a space of automorphic forms.

It should be possible, although I have not done so, to test these suggestions for groups whose unitary representations are well understood, in particular, for $SL(2, R)$ and the De Sitter group. To do this one might make use of an idea basic to Kostant's proof of the (generalized) Borel-Weil theorem for compact groups. Suppose σ is a unitary representation of G on a Hilbert space V . Let $C^q(V)$ be the space of all linear maps from $\wedge^q \mathfrak{n}_\mathbb{C}^-$ to V . $C^q(V)$ is a Hilbert space. The usual coboundary operator from $C^q(V)$ to $C^{q+1}(V)$ can be defined on those elements of $C^q(V)$ which take values in the Gårding subspace of V . The closure d of this operator is the adjoint of the restriction of its formal adjoint to those elements of $C^{q+1}(V)$ which take values in the Gårding subspace. T of course acts on $\wedge^q \mathfrak{n}_\mathbb{C}^-$. If $f \in C^q(V)$ define $tf = f'$ by $f'(X) = tf(t^{-1}X)$, $X \in \wedge^q \mathfrak{n}_\mathbb{C}^-$. There is a natural identification of $C^q(\Lambda)$ with the set of f in $C^q(L^2(G))$ such that $tf = \exp(-\Lambda(H))f$ if $t = \exp H$ belongs to T and of $C^q(\Lambda, \Gamma)$ with the set of f in $C^q(L^2(\Gamma \backslash G))$ such that $tf = \exp(-\Lambda(H))f$. Moreover the following diagrams are commutative.

$$\begin{array}{ccccccc} C^q(\Lambda) & \xrightarrow{\bar{d}} & C^{q+1}(\Lambda) & & C^q(\Lambda, \Gamma) & \xrightarrow{\bar{d}} & C^{q+1}(\Lambda, \Gamma) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ C^q(L^2(G)) & \xrightarrow{d} & C^{q+1}(L^2(G)) & & C^q(L^2(\Gamma \backslash G)) & \xrightarrow{d} & C^{q+1}(L^2(\Gamma \backslash G)). \end{array}$$

The point is that d is easier to study than \bar{d} because to study d we can decompose V into irreducible representations and study the action of d on each part.

References

1. J. Dixmier, *Représentations intégrables du groupe de De Sitter*, Bull. Soc. Math., France **89** (1961), 9-41.
2. Harish-Chandra, *Representations of semisimple Lie groups*, VI, Amer. J. Math. **78** (1956), 564-628.
3. ———, *Discrete series for semisimple Lie groups*, II. Acta. Math. (to appear).
4. Bertram Kostant, *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. (2) **74** (1961), 329-387.
5. R.P. Langlands, *The dimension of spaces of automorphic forms*, Amer. J. Math. **85** (1963), 99-125.