

Bill,

Since you can answer (one hopes) all questions about the unramified principal series, let me remind you of a question I asked a long time ago.

1. G - Chevalley group over non-archimedean local field
2. $G^{\widehat{0}}$ - connected component of associative group
3. $\widehat{\mathfrak{G}}$ - Lie algebra of $G^{\widehat{0}}$.

Consider pairs $\varphi = \{\widehat{t}, X\}$ up to conjugacy, $\widehat{t} \in G^{\widehat{0}}$, \widehat{t} semi-simple. $X \in \widehat{\mathfrak{G}}$, X unipotent. $\text{Ad } \widehat{t}(X) = qX$, q is the number of elements in the residue field.

Can one associate to each such pair φ a finite non-empty set Π_{φ} of irreducible representations of G , which occur as constituents of the unramified principal series such that

- (i) The sets Π_{φ} are disjoint and exhaust the irreducible representations occurring in the principal series
- (ii) The sets Π_{φ} are square integrable \iff there is no proper parabolic subgroup whose Levi factor M contains t while its Lie algebra \mathcal{M} contains X . (A generalization of indecomposable representation.)
- (iii) If ξ : taking $T(F)$, the split Cartan subgroup, to \widehat{L} is defined by

$$|\lambda(t)| = g^{(\text{There is perhaps a minus sign here.})\langle \lambda, \widehat{\lambda} \rangle} \quad \widehat{\lambda} = \xi(t)$$

and if χ is the character of $T(F)$ defined by

$$\chi(t) = \widehat{\lambda}(\widehat{t}) \quad \widehat{\lambda} = \xi(t) \quad \varphi = \{\widehat{t}, X\}$$

then each element of Π_{φ} is a constituent of $PS(\chi)$

- (iv) In $SL(2)$, there is such a pair $\widehat{t} = \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}$, $X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. If one imbeds $SL(2)$ in $G^{\widehat{0}}$ as a principal three-dimensional subgroup Π_{φ} should consist of the special representation alone.

Bob

Compiled on November 17, 2025.