

September 22, 1969
New Haven, CT

Dear Harish-Chandra,

I am very pleased that you are willing to come here to speak. I have decided to come to Princeton on Monday, October 7 and to stay over till Tuesday. However I don't want to give a lecture then because Weil and Jacquet are familiar with anything conclusive I did before January and I have not yet finished the things I have been doing since. I have already given you an idea what these things are but I can perhaps be more precise now.

If F is a local field let C_F be the multiplicative group of F and if F is a global field let C_F be the idèle class group of F . As I said before if K/F is normal the Weil group $G_{K/F}$ is an extension of C_F by the Galois group of K/F . If one likes one can take projective limits and get an object called the Weil group of F . If F is global, and $F_{\mathfrak{p}}$ is a completion of F , and $K_{\mathfrak{p}}$ is a completion of K over \mathfrak{p} then there is a map of $G_{K_{\mathfrak{p}}/F_{\mathfrak{p}}}$ into $G_{K/F}$ and every representation of $G_{K/F}$ determines a representation of $G_{K_{\mathfrak{p}}/F_{\mathfrak{p}}}$. A representation ρ is a finite-dimensional representation over \mathbf{C} such that $\rho(g)$ is semi-simple for all g .

For a non-archimedean local field I can attach to every equivalence class ω of representations a function $L(s, \omega)$ in just one way so that the following conditions are satisfied.

(i) If $\omega \sim \chi_F$ is one-dimensional

$$L(s, \omega) = \frac{1}{1 - \chi_F(\pi_F)|\pi_F|^s} \quad \text{if } \chi_F \text{ is trivial on units}$$

$$= 1 \quad \text{if } \chi_F \text{ is not trivial on units.}$$

(ii) $L(s, \omega_1 \oplus \omega_2) = L(s, \omega_1)L(s, \omega_2)$.

(iii) If $\omega_1 \simeq \text{Ind}(G_{K/F}, G_{K/E}, \omega_2)$ then

$$L(s, \omega_1) = L(s, \omega_2).$$

For archimedean fields $L(s, \omega)$ is defined by the following conditions

(i) If $F = \mathbf{R}$ and $\omega \sim \chi_{\mathbf{R}}$ with $\chi_{\mathbf{R}}(x) = (\text{sgn } x)^m |x|^r$, $m = 0$ or 1 , then

$$L(s, \omega) = \pi^{-\frac{1}{2}(s+r+m)} \Gamma\left(\frac{s+r+m}{2}\right).$$

(ii) If $F = \mathbf{C}$ and $\omega \sim \chi_{\mathbf{C}}$ with $\chi_{\mathbf{C}}(z) = |z|^{2r} \frac{z^n \bar{z}^n}{|z|^{m+n}}$, $m+n \geq 0$, $mn = 0$ then

$$L(s, \omega) = 2(2\pi)^{-(s+r+\frac{m+n}{2})} \Gamma\left(s+r+\frac{m+n}{2}\right)$$

(iii) $L(s, \omega_1 \oplus \omega_2) = L(s, \omega_1)L(s, \omega_2)$

(iv) If $\omega_1 \simeq \text{Ind}(G_{K/F}, G_{K/E}, \omega_2)$ then

$$L(s, \omega_1) = L(s, \omega_2).$$

If K is a global field, ω is an equivalence class of representations of the Weil group and $\omega_{\mathfrak{p}}$ the induced equivalence class of the Weil group of $F_{\mathfrak{p}}$ the Artin L -function is

$$L(s, \omega) = \prod_{\mathfrak{p}} L(s, \omega_{\mathfrak{p}})$$

Let me state the theorem which I have been working on since last January and then comment on its relation to the functional equations of the L -functions. By the way, as soon as I understand a lemma of Dwork I should be able to write out the proof of this theorem. I think I have all the parts of the proof except for some easily managed details. If $F = \mathbf{R}$ and $\chi_{\mathbf{R}}(x) = (\text{sgn } x)^m |x|^r$ with $m = 0$ or 1 and $\psi_{\mathbf{R}}$ is the additive character $\psi_{\mathbf{R}}(x) = e^{2\pi i u x}$ I set

$$\Delta(\chi_{\mathbf{R}}, \psi_{\mathbf{R}}) = (i \text{sgn } u)^m |u|^r.$$

If $F = \mathbf{C}$ and $\chi_{\mathbf{C}}(z) = |z|^{2r} \frac{z^m \bar{z}^n}{|z|^{m+n}}$, $m + n \geq 0$, $mn = 0$ and $\psi_{\mathbf{C}}$ is the additive character $\psi_{\mathbf{C}}(z) = e^{4\pi i \text{Re}(wz)}$ I set

$$\Delta(\chi_{\mathbf{C}}, \psi_{\mathbf{C}}) = i^{m+n} \chi_{\mathbf{C}}(w).$$

If F is non-archimedean, if χ_F is a generalized character of C_F with conductor \mathfrak{P}_F^m , if ψ_F is a non-trivial additive character of F and \mathfrak{P}_F^n is the largest ideal on which it vanishes and $O_F \gamma = \mathfrak{P}_F^{m+n}$ I set

$$\Delta(\chi_F, \psi_F) = \chi_F(\gamma) \frac{\int_{U_F} \psi_F\left(\frac{\alpha}{\gamma}\right) \chi_F^{-1}(\alpha) d\alpha}{\left| \int_{U_F} \psi_F\left(\frac{\alpha}{\gamma}\right) \chi_F^{-1}(\alpha) d\alpha \right|}.$$

U_F is the group of units. The right side is independent of γ .

If E is a separable extension of F and ψ_F is given then

$$\psi_{E/F}(X) = \psi_F(\text{Tr}_{E/F} X).$$

Theorem. *Suppose F is a given local field and ψ_F a given non-trivial additive character of F . It is possible in exactly one way to assign to each separable extension E of F a complex number $\rho(E/F, \psi_F)$ and to each equivalence class ω of representations of the Weil group of E a complex number $\epsilon(\omega, \psi_{E/F})$ so that*

- (i) *If $\omega \simeq \chi_E$ then $\epsilon(\omega, \psi_{E/F}) = \Delta(\chi_E, \psi_{E/F})$*
- (ii) *$\epsilon(\omega_1 \oplus \omega_2, \psi_{E/F}) = \epsilon(\omega_1, \psi_{E/F}) \epsilon(\omega_2, \psi_{E/F})$*
- (iii) *If $\omega_1 \simeq \text{Ind}(G_{K/F}, G_{K/E}, \omega_2)$ then*

$$\epsilon(\omega_1, \psi_F) = \rho(E/F, \psi_F)^{\dim \omega_2} \epsilon(\omega_2, \psi_{E/F}).$$

If A_F^s is the generalized character $\alpha \rightarrow |\alpha|_F^s$ set $\epsilon(s, \omega, \psi_F) = \epsilon(A_F^{s-1/2} \otimes \omega, \psi_F)$. If F is a global field and ψ_F a non-trivial character of \mathbf{A}_F/F let $\psi_{F_{\mathfrak{p}}}$ be the restriction of ψ_F to $F_{\mathfrak{p}}$. If ω is an equivalence class of representations of the Weil group of F and

$$\epsilon(s, \omega) = \prod_{\mathfrak{p}} \epsilon(s, \omega_{\mathfrak{p}}, \psi_{F_{\mathfrak{p}}})$$

the functional equation of the L -function can, on the basis of the previous theorem, be shown to be

$$L(s, \omega) = \epsilon(s, \omega) L(1 - s, \tilde{\omega})$$

if $\tilde{\omega}$ is contragredient to ω .

Once I have this theorem I should very quickly be able to derive a consequence of interest for group representations. However I have not yet carried out the computations. After some preliminaries I shall state the consequence.

If F is a non-archimedean local field a two-dimensional equivalence class ω of representations of the Weil group of F will be called special if ω is the direct sum of two one-dimensional representations μ_F and ν_F and $\mu_F\nu_F^{-1} = A_F^1$ or A_F^{-1} . If $F = \mathbf{R}$ a two-dimensional equivalence class ω is special if $\omega \simeq \mu_F \oplus \nu_F$, and $\mu_F(x) = |x|^{s_1} \left(\frac{x}{|x|}\right)^{m_1}$, $\nu_F(x) = |x|^{s_2} \left(\frac{x}{|x|}\right)^{m_2}$ and $(s_1 - s_2) - (m_1 - m_2)$ is an odd integer. If $F = \mathbf{C}$, ω is special if $\omega \simeq \mu_F \oplus \nu_F$

$$\begin{aligned}\mu_F(z) &= |z|^{2s_1} \left(\frac{z}{|z|}\right)^{m_1} \\ \nu_F(z) &= |z|^{2s_2} \left(\frac{z}{|z|}\right)^{m_2}\end{aligned}$$

and one of $\frac{s_1 - s_2}{2} - \left(1 + \frac{|m_1 - m_2|}{2}\right)$ and $\frac{s_2 - s_1}{2} - \left(1 + \frac{|m_1 - m_2|}{2}\right)$ is a non-negative integer.

$L(\psi_F)$ is the space of functions on $\mathrm{GL}(2, F)$ satisfying

$$\varphi\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}g\right) \equiv \psi_F(x)\varphi(g)$$

together with certain conditions on smoothness and rate of growth. Here is the consequence I mentioned.

Theorem. *Suppose that for every global field F and every two-dimensional irreducible equivalence class of representations of the Weil group of F the function $L(s, \omega)$ is entire and bounded in vertical strips. Then if F is a local field, ω a two-dimensional equivalence class of representations of the Weil group of F which is not special, and ψ_F a non-trivial additive character of F there is a unique simple representation π_ω of $\mathrm{GL}(2, F)$ satisfying*

- (i) π_ω acts on $L \subseteq L(\psi_F)$
- (ii) If φ belongs to L and χ_F is a generalized character of C_F the integral

$$\int_{C_F} \varphi\left(\begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}g\right) \chi_F(\alpha) |\alpha|_F^s d\alpha$$

converges for $\mathrm{Re} s$ sufficiently large. Denote its value by $\Phi(g, s, \varphi, \chi_F)$ and set

$$\Phi(g, s, \varphi, \chi_F) = L\left(s + \frac{1}{2}, \omega \otimes \chi_F\right) \Phi'(g, s, \varphi, \chi_F).$$

$\Phi'(g, s, \varphi, \chi_F)$ is an entire function of s bounded in vertical strips. Moreover

$$\Phi'\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}g, -s, \varphi(\det \omega \chi_F)^{-1}\right) = \epsilon(\chi_F A_F^s \otimes \omega, \psi_F) \Phi'(g, s, \varphi, \chi_F)$$

if $\det \omega$ is the 1-dimensional representation obtained from ω by taking determinants.

It will follow that π_{ω_1} equivalent to π_{ω_2} implies $\omega_1 = \omega_2$. This theorem makes the existence of the representations I mentioned to you earlier pretty much certain. If they do not exist the L -series behave in an entirely unexpected manner.

Yours truly,

Bob Langlands

Compiled on July 3, 2024.